

**MATH 495.595 / PRACTICE FINAL EXAM QUESTIONS
SOLUTIONS / COMMENTS**

1. C is $f(x)$, A is $f'(x)$ and B is $f''(x)$

2. $a + b + c = 99$ $\frac{a}{2} = b$ $a + \frac{a}{2} + c = 99 \Rightarrow c = 99 - \frac{3a}{2}$

Maximize product $a \times b \times c = a \times \frac{a}{2} \times (99 - \frac{3a}{2}) = \frac{99a^2}{2} - \frac{3a^3}{4} = P(a)$ $P'(a) = 99a - \frac{9a^2}{4}$

$0 = 99a - \frac{9a^2}{4} \Rightarrow 99a = \frac{9a^2}{4} \Rightarrow 11 = \frac{a}{4} \Rightarrow a = 44$ Thus, $a = 44$, $b = 22$ and $c = 33$.

3. (Hints) Find x-intercepts, max and min points ($f'(x) = 0$), find where the function is increasing ($f'(x) > 0$), where the function is decreasing ($f'(x) < 0$), where the function is concave up ($f''(x) > 0$), where the function is concave down ($f''(x) < 0$).

4. Suppose a rock is hurled vertically up (on Earth). In the following cases:

a.

(i) Distance: $d(t) = -16t^2 + v_o t + h_o$ feet

Velocity: $v(t) = -32t + v_o$ feet / second

Acceleration: $a(t) = -32$ feet / second²

Given $d(2) = 186$ and $v(3) = 4$

$v(3) = -32(3) + v_o = 4 \Rightarrow v_o = 100 \Rightarrow v(t) = -32t + 100$

$d(t) = -16t^2 + 100t + h_o$

$d(2) = -16(2)^2 + 100(2) + h_o = 186 \Rightarrow h_o = 50 \Rightarrow d(t) = -16t^2 + 100t + 50$

(ii) $d(t) = -16t^2 + 100t + 50 = 0 \Rightarrow t \approx 6.71$, At $v(6.71) = -32(6.71) + 100 \approx -114.7$

The velocity was -114.7 feet / second which means the ball hit the ground, going down, at a speed of 114.7 feet second at $t \approx 6.71$ seconds.

$v(t) = -32t + 100 = 0 \Rightarrow t = 3.125 \Rightarrow d(3.125) = 206.25$

The maximum height of the ball is 206.25 feet above ground

(iii) $v(1.5) = -32(1.5) + 100 = 52$, $v(t) = -32t + 100 = -52 \Rightarrow t = 4.75$

$d(1.5) = 164$ $d(4.75) = 164$

$y - 164 = 52(t - 1.5) \Rightarrow y = 52t + 86$. Tangent line for 1.5 seconds

$y - 164 = -52(t - 4.75) \Rightarrow y = -52t + 411$. Tangent line for 4.75 seconds

(iv) Slope of secant line from any three second interval, example:

$\frac{d(4) - d(1)}{4 - 1} = \frac{60}{3} = 20$ feet / second, average speed of rock from one to four seconds.

b. Proceed in the same way as for part a).

5.


a. $y' = (1 + 3x^2 - x)(1 + 12x^2) + (6x - 1)(x + 4x^3)$

b. $y' = (1 + 7x)13(3x + 4)^{12}(3) + (7)(3x + 4)^{13}$

c. $y' = 12(1 + 7x^3)^{11}(21x^2)$

d. $y' = \frac{(2 - x^4)(21x^2) - (1 + 7x^3)4x^3}{(2 - x^4)^2}$

6. Sketch the function with the following piecewise defined components:



$\lim_{x \rightarrow -2^+} f(x) = 2$

$\lim_{x \rightarrow -2^-} f(x)$; endpoint, can't tell

$\lim_{x \rightarrow 0^+} f(x) = 3$,

$\lim_{x \rightarrow 0} f(x) = DNE$

$\lim_{x \rightarrow 0^-} f(x) = 0$,

$f(-2) = 2$ $f'(x) = -1$, differentiable and continuous

$f(0) = 3$ Jump, not continuous, not differentiable

Limit same as function value, continuous

$\lim_{x \rightarrow 2^+} f(x) = 3$, $\lim_{x \rightarrow 2^-} f(x) = 3$, $\lim_{x \rightarrow 2} f(x) = 3$

$f(2) = 3$

$f'(2) = 0$, from the left

$f'(2) = 2$, from the right

“sharp point” not differentiable

Limit same as function value, continuous

$\lim_{x \rightarrow 5^+} f(x) = 9$, $\lim_{x \rightarrow 5^-} f(x) = 9$, $\lim_{x \rightarrow 5} f(x) = 9$

$f(5) = 9$

$f'(5) = 2$, from the left

$f'(4) = 10$, from the right

“sharp point” not differentiable

$\lim_{x \rightarrow 7^-} f(x) = 33$

$\lim_{x \rightarrow 7^+} f(x)$; endpoint, can't tell

$f(7) = 33$


$f'(x) = 2x$, differentiable and continuous

A function can be continuous and not differentiable at a point (sharp corner), but if a function is not continuous at a point, it is also not differentiable.


7.

a.


i) Derivative




ii) Antiderivatives



iii) Derivative if start second derivative




iv) Function if start second derivative




b.


i) Derivative




ii) Antiderivatives



iii) Derivative if start second derivative




iv) Function if start second derivative




c.


i) Derivative




ii) Antiderivatives



iii) Derivative if start second derivative



iv) Function if start second derivative



8. Using calculus, compute each of the following:

$$\text{a. } \int_{-2}^4 2x^3 + x \, dx = \left. \frac{1}{2}x^4 + \frac{1}{2}x^2 \right]_{-2}^4 = \frac{1}{2}(4)^4 + \frac{1}{2}(4)^2 - \left(\frac{1}{2}(-2)^4 + \frac{1}{2}(-2)^2 \right) = 126$$

$$\text{b. } \int_1^4 \frac{2}{x^3} \, dx = \left. -\frac{1}{x^2} \right]_1^4 = -\frac{1}{(4)^2} - \left(-\frac{1}{1} \right) = .9375$$

$$\text{c. } \int_1^3 \sqrt{2x+1} \, dx = \left. \frac{2}{2 \times 3} \sqrt{(2x+1)^3} \right]_1^3 = \frac{\sqrt{(2(3)+1)^3}}{3} - \frac{\sqrt{(2(1)+1)^3}}{3} = 4.44$$